



## OPTIMIZATION OF SHOOT MULTIPLICATION MEDIA FOR MICRO PROPAGATION OF *WITHANIA SOMNIFERA*: AN ENDANGERED MEDICINAL PLANT

Poonam Duhan<sup>1,2</sup>, Sethi Neeraj<sup>1\*</sup>, Pal Minakshi<sup>2</sup>, Kaura Sushila<sup>3</sup>, Parle Milind<sup>3</sup>

<sup>1</sup>Department of Bio and Nano Technology, Bio and Nano Technology Centre, Guru Jambheshwar University of Science and Technology, Hisar, Haryana, India

<sup>2</sup>Centre for Plant Biotechnology, CCS HAU New Campus, Hisar, Haryana, India

<sup>3</sup>Pharmacology Division, Dept. Pharm. Sciences, Guru Jambheshwar University of Science and Technology, Hisar, Haryana, India

\*Corresponding Author Email: 20neerajsethi@gmail.com

DOI: 10.7897/2277-4572.034168

Published by Moksha Publishing House. Website www.mokshaph.com

All rights reserved.

Received on: 29/05/14 Revised on: 07/07/14 Accepted on: 25/07/14

### ABSTRACT

Effect of different types of plant growth regulators has been studied on *in vitro* multiplication of *Withania somnifera* using nodal explants. Maximum survival percentage 100 % of explants was observed using nodal explants were treated with 0.4 % Streptocyclin and 0.4 % Bavistin for 60 minutes followed by 3 minutes treatment with 0.1 % HgCl<sub>2</sub>. MS medium supplemented with was found to be best for multiplication of nodal explants of *Withania somnifera* amongst seventeen media employed for multiplication. Maximum eight shoots/explants observed on MS basal medium supplemented with 1.5 mg l<sup>-1</sup> BAP + 0.5 mg l<sup>-1</sup> NAA after 45 days of culture. A maximum 77.8 % rooting was found on MS medium (half strength) supplemented with 0.5 mg l<sup>-1</sup> NAA. Shoots with well developed roots were separated individually and transferred to plastic bag containing sand, soil and vermin-compost in ratio of 1:1:1 in the green house and was reported 100 % survival. The hardened plants were successfully transferred to field.

**Keywords:** Micro propagation, Multiplication, Rooting, *Withania somnifera*

### INTRODUCTION

*Withania somnifera*, an herb is one of the common medicinal plants also known as ashwagandha in Sanskrit belongs to family Solanaceae. The plant has been used since Ayurvedic time to treat asthma, bronchitis, emaciation ulcers, insomnia and dementia. It is used as a liver tonic and astringent. Dry arid areas of Bangladesh, China, India and Pakistan are native of this plant. It can be found growing in the Africa and Mediterranean region. The plant possesses many pharmacological actions such as anti-arthritic, anti-rheumatic anti-inflammatory, abortifacient, adaptogenic, anti-stress, anti-tumour, immunomodulatory activities, anti-anxiety, anti-depression and aphrodisiac<sup>1-3</sup>. The medicinal properties of ashwagandha are due to major biochemical constituents such as alkaloids and steroids. Withanolides is main tropane alkaloids of the plant that posses anti-tumour activity. There are few reports available on *in vitro* culture of ashwagandha using different explants, such as from leaf<sup>4-7</sup>, shoot tip<sup>8-10</sup>, axillary buds<sup>11-14</sup>, internodal<sup>15,16</sup>, seed<sup>17</sup> and cotyledon<sup>18,19</sup>. For fulfilling ever increasing demand of pharmaceutical industry for the withanolides and to overcome problem of long gestation period between planting and harvesting of this plant there is great need to produce quality planting material of this plant throughout the year. Micropropagation is a feasible approach for the production of true –to-type planting material. Keeping in view all these points, the present investigation was conducted to study the effect of different plant growth regulators i.e. BAP, KIN and NAA alone and in combination on *in vitro* multiplication of *Withania somnifera* using nodal explants.

### MATERIALS AND METHODS

#### Plant material

Young nodal explants were taken from field grown plant at Medicinal section of CCSHAU, Hisar, Haryana, India.

#### Media Preparation

The present study was conducted using MS<sup>20</sup> basal medium fortified with various plant growth regulators. The pH of medium was adjusted to 5.8 using 0.1N HCL or 0.1N NaOH. Sucrose 30 g l<sup>-1</sup> and agar 8 g l<sup>-1</sup> was added in the medium and autoclaved at 121°C and 15 lbs inch<sup>-2</sup> pressure for 15 minutes.

#### Preparation and sterilization of Explants

Young, healthy nodal segments (2-3 cm) were excised from mother plant and washed with mild detergent for 8-10 minutes followed by rinsing under running tap water to remove detergent. The explants were then giving treatment with 0.4 % Bavistin + 0.4 % Streptocyclin for 60 minutes and followed by washing with double distilled water. Explants were then surface sterilized with 0.1 % HgCl<sub>2</sub> for 1-4 minutes in Laminar air flow and excess HgCl<sub>2</sub> was washed by 4 to 5 washing with double distilled sterilized water.

#### Culture conditions

The cultures vessels were incubated in growth room. The temperature of growth room was maintained at 25 ± 2°C and light intensity of 100 μEM<sup>-2</sup> sec<sup>-1</sup> (1000 lux) was provided using florescent tubes. Period of 16 h / 8 h light and dark was provided.

#### Shoot multiplication

Regenerated shoots were transferred to the MS medium supplemented with various concentrations of BAP (0.5, 1.0, 1.5, 2.0 mg l<sup>-1</sup>), KIN (0.5, 1.0, 1.5, 2.0 mg l<sup>-1</sup>) alone and in combination with NAA (0.5 mg l<sup>-1</sup>) each for multiplication. The data were recorded for number of shoots/explant after 15, 30 and 45 days of inoculation after frequent sub culturing after 2 weeks.

**In vitro rooting**

The well elongated healthy shoots were separated aseptically and transferred individually to the half strength MS medium supplemented with various concentration of auxin NAA (0.5, 1.0, 1.5, 2.0 mg l<sup>-1</sup>).

**Acclimatization and Hardening**

Rooted shoots were transferred in the pots containing different potting mixture of sand, soil and vermin-compost. Pots were kept in the Greenhouse for 4-6 weeks for hardening till the plant attained a height of approximately 15-20 cm. The hardened plants were successfully transferred in the field.

**Table 1: Effect of MS Basal Medium Supplemented with Different Concentrations of Plant Growth Regulators on In Vitro Multiplication of *Withania somnifera***

S. No.	Media code	Average number of shoots/explants after		
		15 day	30 day	45 day
1	WSM0 (MS basal)	0.0 ± 0.00	0.0 ± 0.00	0.0 ± 0.00
2	WSM1 (MS + 0.5 mg l <sup>-1</sup> BAP)	0.8 ± 0.13	1.7 ± 0.08	2.4 ± 0.09
3	WSM2 (MS + 1.0 mg l <sup>-1</sup> BAP)	0.6 ± 0.11	2.2 ± 0.08	2.8 ± 0.11
4	WSM3 (MS + 1.5 mg l <sup>-1</sup> BAP)	1.9 ± 0.27	2.5 ± 0.29	2.5 ± 0.12
5	WSM4 (MS + 2.0 mg l <sup>-1</sup> BAP)	1.7 ± 0.08	2.2 ± 0.08	3.6 ± 0.10
6	WSM5 (MS + 0.5 mg l <sup>-1</sup> KIN)	0.5 ± 0.13	0.5 ± 0.13	0.8 ± 0.13
7	WSM6 (MS + 1.0 mg l <sup>-1</sup> KIN)	1.0 ± 0.10	1.7 ± 0.08	2.5 ± 0.12
8	WSM7 (MS + 1.5 mg l <sup>-1</sup> KIN)	1.0 ± 0.00	2.4 ± 0.09	2.7 ± 0.62
9	WSM8 (MS + 2.0 mg l <sup>-1</sup> KIN)	1.0 ± 0.00	2.4 ± 0.09	2.5 ± 0.12
10	WSM9 (MS + 0.5 mg l <sup>-1</sup> BAP + 0.5 mg l <sup>-1</sup> NAA)	2.1 ± 0.26	3.2 ± 0.56	5.6 ± 0.33
11	WSM10 (MS + 1.0 mg l <sup>-1</sup> BAP + 0.5 mg l <sup>-1</sup> NAA)	3.8 ± 0.20	4.6 ± 0.23	5.8 ± 0.27
12	WSM11 (MS + 1.5 mg l <sup>-1</sup> BAP + 0.5 mg l <sup>-1</sup> NAA)	6.7 ± 0.3	7.6 ± 0.54	8.0 ± 0.57
13	WSM12 (MS + 2.0 mg l <sup>-1</sup> BAP + 0.5 mg l <sup>-1</sup> NAA)	5.8 ± 0.28	6.7 ± 0.3	7.2 ± 0.34
14	WSM13 (MS + 0.5 mg l <sup>-1</sup> KIN + 0.5 mg l <sup>-1</sup> NAA)	2.0 ± 0.22	4.5 ± 0.57	5.1 ± 0.82
15	WSM14 (MS + 1.0 mg l <sup>-1</sup> KIN + 0.5 mg l <sup>-1</sup> NAA)	1.6 ± 0.07	2.5 ± 0.29	4.5 ± 0.57
16	WSM15 (MS + 1.5 mg l <sup>-1</sup> KIN + 0.5 mg l <sup>-1</sup> NAA)	1.6 ± 0.07	3.2 ± 0.56	4.9 ± 0.63
17	WSM16 (MS + 2.0 mg l <sup>-1</sup> KIN + 0.5 mg l <sup>-1</sup> NAA)	2.7 ± 0.62	3.2 ± 0.56	3.2 ± 0.56

\*Data is shown as mean of three replicates ± S.E.

**Table 2: Effect of MS (Half Strength) Basal Medium Supplemented with Different Concentrations (0.5-2.0 mg l<sup>-1</sup>) of Auxin (Naa) on Percentage Root Induction of Plantlets of *Withania somnifera***

S. No.	Media code	% root induction	No. of days required for rooting
1	WSR0 (½ MS Basal)	0.00 ± 0.00	-
2	WSR1 (½ MS + 0.5 mg l <sup>-1</sup> NAA)	77.8 ± 0.08	8.1 ± 0.40
3	WSR2 (½ MS + 1.0 mg l <sup>-1</sup> NAA)	55.6 ± 0.06	13.5 ± 0.38
4	WSR3 (½ MS + 1.5 mg l <sup>-1</sup> NAA)	33.3 ± 0.03	15.3 ± 0.44
5	WSR4 (½ MS + 2.0 mg l <sup>-1</sup> NAA)	22.2 ± 0.02	17.7 ± 0.44

\*Data are shown as mean of three replicates ± S.E. - = no rooting

**Table 3: Effect of Different Potting Mixture on Hardening of In Vitro Raised Plantlets of *Withania somnifera***

S. No.	Mixture Code	Potting mixture	% survival of rooted plantlets
1	WSPM1	Sand + FYM (1:1)	80
2	WSPM2	Sand + Soil + FYM (1:1:1)	80
3	WSPM3	Sand + Soil + vermin compost (1:1:1)	100



**Figure 1: In vitro establishment, multiplication and rooting of *Withania somnifera***

(A) Regeneration of nodal explants on MS basal media, (B) Initiation of proliferation and multiple shoots from regenerated shoots on MS + 1.5 mg l<sup>-1</sup> BAP + 0.5 mg l<sup>-1</sup> NAA after 15 days (C) multiple shoots on MS + 1.5 mg l<sup>-1</sup> BAP + 0.5 mg l<sup>-1</sup> NAA after 45 days (D) In vitro rooting on MS + 0.5 mg l<sup>-1</sup> NAA

## RESULTS AND DISCUSSION

Effect of different duration of sterilization agents Bavistin, Streptocyclin and 0.1 %  $\text{HgCl}_2$  studied on *in vitro* establishment of *Withania somnifera*. The explants were treated with 0.1 %  $\text{HgCl}_2$  for different duration (1-4 minutes) and data were recorded for percent contamination and percent survival after 21 days of inoculation. A maximum 100 % survival and asepsis was observed when nodal explants treated with 0.4 % Streptocyclin and 0.4 % Bavistin for 60 minutes followed by 3 minutes treatment with 0.1 %  $\text{HgCl}_2$ . Long duration of  $\text{HgCl}_2$  exceeding above 3 minutes treatment was resulted in browning and mortality of explants. Similar results were reported by<sup>21</sup> using 0.01 %  $\text{HgCl}_2$  for 3-5 minutes in *Celastrus paniculatus*. Similarly, also found similar result in *Achyranthes aspera* medicinal plant using nodal explants<sup>22</sup>. In the present investigation effect of different types and concentrations of plant growth regulators Viz. BAP (0.5-2.0  $\text{mg l}^{-1}$ ), KIN (0.5-2.0  $\text{mg l}^{-1}$ ) alone and each in combination with (0.1  $\text{mg l}^{-1}$ ) NAA was employed for *in vitro* multiplication of nodal explants of *Withania somnifera* and data were recorded for number of shoots/explants on fixed intervals of time (Table 1). Seventeen different media combinations (WSM<sub>0</sub> -WSM<sub>16</sub>) employed for multiplication of nodal explants, MS medium supplemented with 1.5  $\text{mg l}^{-1}$  BAP + 0.1  $\text{mg l}^{-1}$  NAA was found to be best for multiplication. The maximum shoots/explant (8.0) was observed on medium WSM<sub>11</sub> (MS + 1.5  $\text{mg l}^{-1}$  BAP + 0.5  $\text{mg l}^{-1}$  NAA) followed by medium WSM<sub>12</sub> (MS + 2.0  $\text{mg l}^{-1}$  BAP + 0.5  $\text{mg l}^{-1}$  NAA) with 7.2 shoots/explants after 45 days (Table 1; Figure 1.C). Minimum shoots (0.8) were observed on WSM<sub>5</sub> medium MS basal medium supplemented with 0.5  $\text{mg l}^{-1}$  KIN. It was observed that BAP was more effective than KIN for shoot multiplication (Figure 1 B, C). BAP was the most effective cytokinin for promoting shoot multiplication and proliferation from nodal segments. Results of present study are in close agreement with earlier reports for shoot multiplication in ashwagandha by several researchers<sup>13, 18,23</sup> suggesting BAP was most effective cytokinin. In the present study the lower concentration of cytokinin (BAP) in combination with lower auxin (NAA) was found to be superior over KIN alone and in combination with NAA for shoot proliferation of *Withania somnifera*. Result of present study in close agreement with<sup>24</sup> reported that BAP with NAA in low concentration produced direct shoots with a little callus development. While, 13 reported combination of 1.0 BAP and 1.0 KIN best for shoot multiplication in ashwagandha using axillary explants. Hence the data presented here supports that BAP is better than KIN for *in vitro* multiplication of *Withania somnifera*. For root induction, the well developed shoots were separated aseptically and were transferred to the half strength MS media supplemented with various concentration of IAA (0.5-2.0  $\text{mg l}^{-1}$ ). The highest rooting (77.8 %) was observed within minimum 8.1 days on the WSR<sub>1</sub> medium supplemented with MS medium (half strength) medium supplemented with 0.5  $\text{mg l}^{-1}$  NAA followed by medium WSR<sub>2</sub> (½ MS + 1.0  $\text{mg l}^{-1}$  NAA) with 55.6 % rooting (Table 2). In present it was observed that on increasing concentration of NAA above 0.5  $\text{mg l}^{-1}$  result in decrease rooting percentage. Low concentration of NAA was found more effective for root induction in *Withania somnifera* (Figure 1 D). Contrary, 85-90 % rooting using ms medium supplemented with 200  $\text{mg l}^{-1}$  activated charcoal within 15-20 days<sup>24</sup>. Therefore, the data support that it is better to use lower concentration of auxins

for *in vitro* root induction of *Withania somnifera*. Rooted shoots were separated individually and transferred to plastic bags containing different potting mixture of sand, soil and vermin-compost. A maximum 100 % survival was reported on potting mixture WSPM<sub>4</sub> (sand, soil and vermin-compost in ratio of 1:1:1). After that hardened plants transferred to pot or in field for commercial purpose (Table 3)

## CONCLUSION

Amongst the different media employed MS media was found to be most suitable for growth and development of *Withania somnifera*. A high frequency of multiple shoots were obtained on MS media supplemented with 1.5  $\text{mg l}^{-1}$  BAP supplemented with 0.5  $\text{mg l}^{-1}$  NAA. In addition, the half strength MS medium supplemented with 0.5  $\text{mg l}^{-1}$  NAA proved to be most effective for root induction. Thus in future this protocol will serve very useful for *in vitro* propagation of *Withania somnifera*.

## REFERENCES

1. Devi PU, Sharada AC. *In vivo* growth inhibitory effect of *Withania somnifera* (Ashwaganda) on a transplantable mouse tumor, Sarcoma 180. India Journal of Experimental Biology 1992; 30(3): 169-17.
2. Devi PU, Sharada AC, Solomon SE. *In vivo* growth inhibitory and radio sensitivity affects of withaferin A, on mouse. Cancer Lett 1995; 95: 189-193. [http://dx.doi.org/10.1016/0304-3835\(95\)03892-Z](http://dx.doi.org/10.1016/0304-3835(95)03892-Z)
3. Devi PU. *Withania somnifera* Dunal (Ashwaganda): Potential plant source of a promising drug for cancer chemotherapy and radio sensitization. India Journal of Experimental Biology 1996; 34(10): 927-932.
4. Baburaj S, Gunakeswaran K. *In vitro* differentiation of shoots from leaf callus culture of *Withania somnifera* (L.) dunal. J. Indian Bot. Soc 1995; 74: 323-324.
5. Kulkarni A, Thengane SR, Krishnamurthy KV. Direct *in vitro* regeneration of leaf explant of *Withania somnifera* (L.) dunal. Plant Science 1996; 119: 163-168. [http://dx.doi.org/10.1016/0168-9452\(96\)04462-7](http://dx.doi.org/10.1016/0168-9452(96)04462-7)
6. Sivanesan I and Murugesan K. *In vitro* adventitious shoot formation from leaf explants of *Withania somnifera* Dunal. Plant Cell Biotechnology and Molecular Biology 2005; 6 (3, 4): 163-166.
7. Joshi AG, Padhya MA. Shoot regeneration from leaf explants of *Withania somnifera* (L.) Dunal. Notulae Scientia Biologicae 2010; 2(1): 63-65.
8. Sen J, Sharma AK. Micropropagation of *Withania somnifera* from germinating seeds and shoot tips. Plant Cell, Tissue and Organ Culture 1991; 26: 71-73. <http://dx.doi.org/10.1007/BF00036108>
9. Sivanesan I. Direct regeneration from apical bud explants of *Withania somnifera* Dunal. Indian Journal of Biotechnology 2007; 6: 125-127.
10. De Silva MAN, Senarath WTPSK. *In vitro* mass propagation and greenhouse establishment of *Withania somnifera* (L.) Dunal (Solanaceae) and comparison of growth and chemical compounds of tissue cultured and seed raised plants. Journal of the National Science Foundation of Sri Lanka 2009; 37(4): 249-255.
11. Roja G, Heble MR. Tissue culture of *Withania somnifera*: Morphogenesis and withanolide synthesis. Phytotherapy Research 1991; 5: 185-187. <http://dx.doi.org/10.1002/ptr.2650050411>
12. Gita R, Grover IS. *In vitro* callus induction and regeneration studies in *Withania somnifera*. Plant Cell, Tissue and Organ Culture 1999; 57(1): 23-27. <http://dx.doi.org/10.1023/A:1006329532561>
13. Sabir F, Sangwan NS, Chaurasiya ND, Misra LN, Tuli R, Sangwan RS. Rapid micropropagation of *Withania somnifera* L. accessions from axillary meristems. Journal of Herbs, Spices and Medicinal Plants 2008; 13(4): 123-132. <http://dx.doi.org/10.1080/10496470801946166>
14. Valizadeh J and Valizadeh M. Development of efficient micropropagation protocol for *Withania coagulans* (Stocks) Dunal. African Journal of Biotechnology 2011; 10 (39): 7611-7616.
15. Govindaraju B, Rao SR, Venugopal RB, Kiran SG, Kaviraj CP, Rao S. High frequency plant regeneration in Ashwaganda (*Withania somnifera* (L.) Dunal): An important medicinal plant. Plant Cell Biotechnology and Molecular Biology 2003; 4(1): 49-56.
16. Valizadeh J, Valizadeh M. *In vitro* callus induction and plant regeneration from *Withania coagulans*: A valuable medicinal plant. Pakistan Journal of Biological Sciences 2009; 12(21): 1415-1419. <http://dx.doi.org/10.3923/pjbs.2009.1415.1419>
17. Supe U, Dhote F, Roymon MG. *In Vitro* plant regeneration of *Withania somnifera*. Plant Tissue Cult. and Biotech 2006; 16(2): 11-115.

18. Rani GG, Virk GS, Nagpal A. Callus induction and plantlet regeneration in *Withania somnifera* (L.) dunal. *In Vitro Cellular and Developmental Biology - Plant* 2003; 39: 468–474. <http://dx.doi.org/10.1079/IVP2003449>
19. Jhankare A, Tiwari G, Tripathi MK, Baghel BS, Tiwari S. Plant regeneration from mature cotyledon, embryo and hypocotyls explants of *Withania somnifera* (L.) Dunal. *Journal of Agricultural Technology* 2011; 7(4): 1023-1035.
20. Murashige T, Skoog F. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol Plant* 1962; 15: 473-497. <http://dx.doi.org/10.1111/j.1399-3054.1962.tb08052.x>
21. Lal D, Singh N. Mass multiplication of *Celastrus paniculatus* Willd: An important medicinal plant under *in vitro* conditions via nodal segments. *International Journal of Biodiversity and Conservation* 2010; 2(6): 140-145.
22. Sen MK, Dash BK, Jerin RS, Faruquee HM. Phytochemistry and antimicrobial potential of apang (*Achyranthes aspera* L.): An update review. *International Journal of Pharmaceutical Sciences Review and Research* 2012; 3: 1-8.
23. Ray S, Jha S. Production of withaferin A in shoot culture of *Withania somnifera*. *Planta* 2001; 67: 432-436.
24. Baba IA, Alia A, Saxena RC, Itoo A, Kumar S, Ahmad M. *In vitro* propagation of *Withania Somnifera* (L.) Dunal (Ashwagandha) an endangered medicinal plant. *International Journal of Pharmaceutical Science Invention* 2013; 2(3): 6-11.

Source of support: Nil, Conflict of interest: None Declared

<b>QUICK RESPONSE CODE</b> 	ISSN (Online) : 2277 –4572
	Website <a href="http://www.jpsionline.com">http://www.jpsionline.com</a>

**How to cite this article:**

Poonam Duhan, Sethi Neeraj, Pal Minakshi, Kaura Sushila, Parle Milind. Optimization of shoot multiplication media for micro propagation of *Withania somnifera*: An endangered medicinal plant. *J Pharm Sci Innov.* 2014;3(4):340-343 <http://dx.doi.org/10.7897/2277-4572.034168>